



FUNGI-PHITE®
EPA REG. NO.: 73771-5

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label. This recommendation must be in the possession of the user at the time of pesticide application.

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

PECANS

DISEASE	APPLICATION METHOD	APPLICATION RATE	APPLICATION PROGRAM*
Scab (<i>Fusicladium effusum</i>)	Foliar Spray	Aerial: Apply 2-4 quarts/acre (5-9 L/ha) in a minimum of 20 gallons/acre (190 L/ha) of water.	Apply in a preventative spray program beginning at bud break, directing the spray to all parts of the tree. Continue applications at 7-10-day intervals. Use a tank mix partner registered for scab or downy spot control in alternate sprays.
Downy spot (<i>Mycosphaerella caryigena</i>)		Ground: Apply 2-4 quarts/acre (5-9 L/ha) in a minimum of 100 gallons/acre (950 L/ha) of water.	Do not apply after shuck split or within 30 days of harvest. Do not apply more than 6 times per crop cycle.

* Check with your local Extension Agent or Crop Consultant if you are unsure about disease prevention, control or severity/pressure.

Read the “Conditions of Sale and Limitations of Warranty and Liability” in the label booklet for Fungi-Phite before using.