

INTRACEPT

F002331

GUARANTEED ANALYSIS

Sulfur (S)	1.00%
Copper (Cu)	0.10%
0.10% Complexed Copper	
Iron (Fe)	0.10%
0.10% Complexed Iron	
Manganese (Mn)	0.33%
0.33% Complexed Manganese	
Zinc (Zn)	0.40%
0.40% Complexed Zinc	

Derived from copper amino acid complex, iron amino acid complex, manganese amino acid complex, and zinc amino acid complex.



ANGER

Causes serious eye damage. May cause damage to organs through prolonged or repeated exposure.

PRECAUTIONARY STATEMENTS

PREVENTION

Wear eye protection/ face protection. Do not breathe dust/fume/gas/mist/vapors/spray.

RESPONSE

Get medical advice/attention if you feel unwell.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or doctor.

Causes mild skin irritation. Very toxic to aquatic life with long lasting effects.

USE PRECAUTIONS - KEEP OUT OF REACH OF CHILDREN

As with all materials, user should exercise precaution to avoid potential phytotoxicity including foliar burning. Care is advised if applying this product in combination with other materials, a jar test for compatibility is recommended before tank mixing with other products. User assumes full responsibility to ensure product, equipment and plant compatibilities. Personal Protective Equipment. Intended for Agricultural use. Consult Safety Data Sheet for further information. In the event of a spill, clean-up should be done promptly by trained personnel while using pre-planned procedures.

GENERAL RECOMMENDATIONS

Intracept is a complex micronutrient designated to be foliar or soil applied. It has been developed to enhance nutrient penetration, uptake and utilization. It is readily available and rapidly absorbed by the plant, due to its complexing properties. This product may be used on a variety of crops and should be applied at planting or at the beginning of stressful growth stages such as first true leaf, flowering and fruit set. It's water soluble and non-toxic to plants when applied as directed.

IRRIGATION SYSTEM APPLICATION

Applications can be made through certain irrigation systems including, but not limited to, low volume systems and typical agricultural sprinkler systems. The specific injection equipment and irrigation system must conform to all federal, state and local regulations pertaining to the fertilization of plants through an irrigation system. Accurate and properly calibrated injection equipment and an irrigation system that is operated at maximum distribution uniformity will help avoid plant injury and/or product ineffectiveness. For best results, with the exception of center pivot and lateral systems, apply this product toward the end of the irrigation set. Allow sufficient time for the product to be flushed completely out of the entire irrigation system prior to turning off the irrigation water.

STORAGE AND DISPOSAL

Store in a cool, dry place. Keep container tightly closed and out of direct sunlight during storage for extended amounts of time. Do not add water or other material to the container. Do not contaminate water, food, or feed by storage and disposal. Store above 32°F. In case of a spill, clean up promptly using trained personnel and pre-planned procedures. Dispose of content and/or container in accordance with local, regional, national and/or international regulations.

DIRECTIONS FOR USE

Foliar Application: Apply in an adequate amount of water that will provide complete coverage of the plants. Use of a non ionic wetting agent may improve spray coverage of certain hard to wet plants. This product may be included in a regular spray program on crops.

Soil Application: In a planter band application a soil barrier between the seed and the product is preferred.

Contact your Crop Advisor for specific recommendations for your area.

Potatoes: Apply 16 to 32 fl. oz. per acre. May be applied in a planter band followed by a foliar application at the appropriate time. Applications may be repeated two or more times throughout the growing season.

Fruit Trees: Apply 16 to 32 fl. oz. per acre before blossoming and shortly after petal fall.

Melons and Vine Crops: Apply 16 to 32 fl. oz. per acre at first female flower and again at softball-sized fruit.

Onions: Apply 16 to 32 fl. oz. per acre at bulb initiation (1/2" to 3/4" diameter). A second application (1.5" to 2" bulb diameter) is recommended. Applications may be repeated two or more times throughout the growing season. Observe all precautions and limitations on this label and labels of products used with Intracept.

Lot No.: XXXXXX

Mfg Date: MM/DD/YYYY

Exp Date: MM/DD/YYYY

Net Contents: 2.5 gal (9.46 L)

Net Wt.: 24.38 lbs. (11.1 kg)

Inv. ID: 1000546

Density: 9.75 lbs./gal (1.17 kg/L) at 68 °F (20 °C)

In case of emergency call INFOTRAC: 800-535-5053 or INTL INFOTRAC: 352-323-3500

To access Verdesian's Warranty Statement please visit <https://www.vlsci.com/limited-warranty>

Contents and levels of metals information for this product is available at <https://www.aapco.org/metals.html>



VERDESIAN

THE NUTRIENT USE EFFICIENCY PEOPLE®

This product is guaranteed by Verdesian Life Sciences, U.S., LLC

1001 Winstead Drive, Suite 480, Cary, NC 27513

Customer Service: 800-868-6446

© 2025 Verdesian Life Sciences U.S., LLC. All rights reserved.

A VERDESIAN NUE SOLUTION®
INTRACEPT