

TAKE OFF ST®

Seed Treatment for Vegetable, Field, and Permanent Crops

F002331

COMPOSITION

CONTAINS NON-PLANT FOOD INGREDIENTS

Active Ingredients:

Citric Acid	0.75%
Glutamate	0.25%
Proline	0.25%
Inert Ingredients (Water)	98.75%

This product is not considered hazardous by the US OSHA Hazard Communication Standard 2024 (29 CFR 1910.1200).

USE PRECAUTIONS - KEEP OUT OF REACH OF CHILDREN

Note: Tank mix environments where the pH is less than 2.5 reduce the efficacy of the product and as such should be avoided.

As with all materials, user should exercise precaution to avoid potential phytotoxicity including foliar burning. Caution is advised if applying this product in combination with other materials, a jar test for compatibility is recommended before tank mixing with other products. User assumes full responsibility to ensure product, equipment and plant compatibilities. Personal Protective Equipment. Intended for Agricultural use. Consult Safety Data Sheet for further information. In the event of a spill, clean-up should be done promptly by trained personnel while using pre-planned procedures.

GENERAL RECOMMENDATIONS

Take Off® ST is a proprietary formulation and blend of organic acid and amino acids for use as a seed treatment on a wide range of plant seeds including, but not limited to, cereals and vegetables. This product has seed priming properties that are beneficial when applied as a seed treatment.

This product is not recommended for use as a fertilizer substitute.

Contact your Crop Advisor for specific recommendations for your area.

STORAGE AND DISPOSAL

Store in a cool, dry area out of reach of children and animals, as well as out of direct sunlight during storage for extended amounts of time. Store in original container. Keep container tightly sealed when not in use. Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Store at temperatures above 32°F. Dispose of unused product and empty containers at an approved waste disposal plant in accordance with Federal, State and local regulations.

DIRECTIONS FOR USE

This product can be applied to seeds with most seed treating equipment commonly used in the industry. Follow seed application recommendations listed below for specific seed size/type.

SEED APPLICATION RECOMMENDATIONS

Recommendation by Specific Crop	Application Rate	Minimum Dilution per 100# of seed
barley, rice, rye, wheat, cotton, canola, sugar beets	0.3 oz/100#	3 fl oz
soybeans	0.3 oz/140,000 seeds or 0.0678 mg/seed	6 fl oz
corn	0.6 oz/80,000 or 0.2373 mg/seed	6 fl oz

Example: Take Off ST Application Rate = 0.3 oz diluted in 3 fl oz of water per 100# of seed.

General Recommendation by seed size	Application Rate/Ton	Minimum Dilution Rate/Ton of seed
Very Small	3 oz (100 ml)	1 qt (1 ltr)
Small	6 oz (200 ml)	2 qt (2 ltr)
Medium	9 oz (300 ml)	3 qt (3 ltr)
Large	12 oz (400 ml)	4 qt (4 ltr)

Example: Take Off ST Application Rate/Ton for wheat = 0.3 oz diluted in 3 oz of water per 100# of seed.

Compatibility – This product is compatible with most common foliar pesticides and nutrients. It is recommended that a small scale compatibility test be conducted prior to mixing a full batch.

A VERDESIAN NUE SOLUTION™

TAKE OFF ST®

To access Verdesian's Warranty Statement please visit vlsci.com/limited-warranty.

Contents and levels of metals information for this product is available at aapfco.org/metals.html.

In case of emergency call INFOTRAC: 800-535-5053 or INTL INFOTRAC: 352-323-3500

Lot No.: XXXXXX

Mfg Date: MM/DD/YYYY

Contents: 2.5 gal (9.46 L)

Exp Date: MM/DD/YYYY

Net Wt.: 22.30 lbs. (10.12 kg)

Density: 8.92 lbs./gal (1.07 kg/L) at 68° F (20° C)

Inv. ID: 1000739



VERDESIAN

THE NUTRIENT USE EFFICIENCY PEOPLE®

This product is guaranteed by Verdesian Life Sciences, U.S., LLC

1001 Winstead Drive, Suite 480, Cary, NC 27513 | Customer Service: 800-868-6446

© 2025 Verdesian Life Sciences U.S., LLC. All rights reserved.